

# Density fluctuations in one-field inflation

Masahide Yamaguchi

*Department of Physics and Mathematics,  
Aoyama Gakuin University, Kanagawa 229-8558, Japan*

Jun'ichi Yokoyama

*Research Center for the Early Universe (RESCEU),  
Graduate School of Science, The University of Tokyo, Tokyo, 113-0033, Japan*

(Dated: February 2, 2008)

## Abstract

Any one-field inflation is actually realized in a multifield configuration because the inflaton must have couplings with other fields to reheat the universe and is coupled to all other fields at least gravitationally. In all single inflaton models, it is explicitly or implicitly assumed that the heavier fields are stuck to their potential minima during inflation, which are time-dependent in general. We present a formalism to calculate curvature perturbations in such a time-dependent background and show that the proper expression can be obtained using a single-field analysis with a reduced potential in which all these heavy fields are situated at their respective, time-dependent minima. Our results provide a firm ground on the conventional calculation.

PACS numbers: 98.80.Cq

We live in a big, spatially flat, globally homogeneous and isotropic universe full of entropy today— all by virtue of inflation in the early universe [1]. The slow-roll inflation also generates primordial density fluctuations as a seed for the large scale structure of the universe [2]. Recent observations of the cosmic microwave background fluctuations by the WMAP [3, 4] and the BOOMERANG-03 [5] found the concordance with the predictions of inflation and confirmed that primordial density fluctuations are adiabatic, Gaussian, and nearly scale-invariant. Furthermore, the observation of the correlation between the temperature anisotropy and the E-mode polarization has confirmed inflation-produced adiabatic fluctuations and falsified causal seed models and primordially isocurvature fluctuations as a generation mechanism of curvature perturbation [4].

The simplest models of inflation require only one real scalar field, which we call the *inflaton*, and inflation is driven as it slowly rolls down along its potential toward a minimum. The inflaton, however, is by no means the sole scalar field of the theory because the inflaton must have couplings with other fields to reheat the universe and is coupled to all other fields at least gravitationally. Furthermore, a lot of inflation models are recently considered in the stringy landscape [6], in which there are a large number of scalar fields with a huge number of extrema. We simply ignore the existence of the other fields by postulating either that they are anchored at their respective minima during inflation due to some potential, or that their energy scale is so low that neither their homogeneous parts nor their fluctuations play important roles. Thus we should always keep in mind that inflation occurs in the multifield configuration in a sensible theory of high energy physics.

A number of analyses have been done on inflation models with multiple scalar fields, which involve many fluctuating fields, focused on the generation of isocurvature fluctuations in addition to the adiabatic fluctuations [7]. To our knowledge, however, the other important class of inflation models with multiple scalar fields mentioned above has not been fully investigated yet, in which only one field  $\varphi$  is light and the others  $\chi_i$  ( $i = 1, \dots, N$ ) are heavy. In such a situation, the light field  $\varphi$  plays a role of the inflaton but the other heavy fields  $\chi_i$  may affect the trajectory of the light field. In fact, virtually all the single inflaton models based on modern unified theories fall into this category for the reason stated above, either manifestly or implicitly. In particular, inflation models are often considered in the stringy landscape, in which there are a large number of scalar fields with a huge number of extrema. In the folded inflation model, for example, the inflaton turns several corners in the potential [8].

Among the manifestly multifield models, the best known example is the hybrid inflation model proposed by Linde [9], which is recently paid particular attention to as a model to realize the running feature of the spectral index of density fluctuations inferred by the WMAP results [10]. In hybrid inflation, the heavy fields are stuck on the origin  $\chi_i = 0$  due to their heavy masses during inflation so that the light field  $\varphi$  rolls down along the effective potential  $V(\varphi) = V(\varphi, \chi_i = 0)$ . Thus, we look on hybrid inflation as an effectively single field inflation and calculate primordial density perturbations by use of the effective potential  $V(\varphi) = V(\varphi, \chi_i = 0)$ . The original hybrid inflation, however, is exceptional among the manifestly multifield models because the light and the heavy fields are orthogonal to each other in the sense that the minima  $\chi_i^m = 0$  are independent of the light field.

We must point out, however, that in general the minima of heavy fields are dependent on the inflaton. This is because all the scalar fields are coupled with each other at least with the gravitational strength. Then the expectation values of other heavy fields may well shift significantly during inflation, being affected by the motion of the inflaton. For example, the

number of  $e$ -folds elapsed while the inflaton changes by  $\Delta\phi$  is given by

$$N = 8\pi G \left| \int_{\phi-\Delta\phi}^{\phi} \frac{V(\phi)}{V'(\phi)} d\phi \right| \approx \frac{(\Delta\phi)^2}{M_G^2}, \quad (1)$$

for a typical polynomial type potential of the inflaton  $V(\phi)$  which is used to realize chaotic inflation [11]. Thus during the last 55  $e$ -folds corresponding to the observable regime, the inflaton changes more than the reduced Planck scale  $M_G = 2.4 \times 10^{18} \text{GeV}$ . This is also the case with many small-field models such as new [12] and topological inflation models [13] because the vacuum expectation value after inflation typically takes a value of order of  $M_G$ . Therefore, this change would inevitably affect the behavior of other heavy fields even if they are coupled with the inflaton only through gravitational strength, whose back reaction to the inflaton may not be negligible in turn.

Furthermore in some types of small-field inflation models, in which coupling constants of interactions of heavy fields are much less than unity, the potential minima of these heavy fields can be modulated considerably thorough gravitationally suppressed interactions, even when change of the inflaton is much *smaller* than  $M_G$ . A good example is the smooth hybrid inflation proposed by Lazarides and Panagiotakopoulos [14]. In this model, the potential for a light field  $\varphi$ , which plays the role of the inflaton, and a heavy field  $\chi$  is given by

$$V(\varphi, \chi) = \left( \mu^2 - \frac{\chi^4}{16M^2} \right)^2 + \frac{\chi^6 \varphi^2}{16M^4}, \quad (2)$$

where  $\mu$  and  $M$  are some mass scales. For  $\varphi^2 \gg \mu M$ , the potential minimum of  $\chi$  is given by

$$(\chi^m)^2 \simeq \frac{4}{3} \frac{\mu^2 M^2}{\varphi^2}. \quad (3)$$

Note that  $\chi^m$  is dependent on  $\varphi$  here, that is,  $\chi^m = \chi^m(\varphi)$ . Since the effective mass squared at the minimum  $\chi^m$ ,  $m_\chi^2 \simeq 4\mu^4/(3\varphi^2)$ , is much larger than the Hubble parameter squared,  $H^2 \simeq \mu^4/(3M_G^2)$ , as long as  $\varphi \ll M_G$ ,  $\chi$  quickly relaxes at the minimum during inflation. Then, the dynamics of the homogeneous mode of the  $\varphi$  field is described by the effective single-field potential  $V_s(\varphi) = V(\varphi, \chi = \chi^m(\varphi))$  and leads to inflation. Recently, this type of inflation is found to be attractive [15, 16] because, combined with new inflation [12], it can simultaneously explain the two new features discovered by the recent precision measurements of cosmic microwave background anisotropy, that is, the running of spectral index of density fluctuations on large scale as preferred by the first-year WMAP data and the BOOMERANG-03, and a large enough amplitude of fluctuation on small scale relevant to first star formation to realize early re-ionization as discovered by WMAP.

In [15], however, primordial density fluctuations have been calculated by looking on such models as effectively single-field inflation of the inflaton  $\varphi$  with the potential  $V_s(\varphi) = V(\varphi, \chi = \chi^m(\varphi))$ . Although it is true that the dynamics of the homogeneous mode is determined by the effective single-field potential, it is a nontrivial task to justify the above procedure to calculate primordial density fluctuations, which enables us to estimate the effects of the change of the minima of heavy fields on density fluctuations for the first time.

As mentioned before, the existence of the other fields is often ignored simply postulating either that they are so heavy that they are anchored at their respective minima during inflation, or that their energy scale is so low that neither their homogeneous parts nor their fluctuations play important roles. However, the considerations about the mass scale or the

energy scale of heavy fields are not sufficient to estimate their effects on density fluctuations. That is, even if changes of the minima of heavy fields are very small, there is no guarantee that such changes do not have significant effects on density fluctuations. Indeed they depend not only on the potential energy density, which is hardly affected by the small change of heavy fields, but also the kinetic term, for their amplitude is inversely proportional to the time derivative of the inflaton. Since its kinetic energy is much smaller than the potential energy during inflation, a small shift in the heavy field may well induce a large relative change in the kinetic term, which may affect the amplitude of density fluctuation considerably.

In this paper, we investigate this problem in detail. That is, we consider inflation models with multiple scalar fields, in which only one field  $\varphi$  is light and the others  $\chi_i$  ( $i = 1, \dots, N$ ) are heavy. During inflation, the heavy fields  $\chi_i$  are stuck to their minima  $\chi_i^m(\varphi)$ , which can depend on the light field  $\varphi$ . We study creation of perturbations in this setting. Since there is only one light mode that acquires long-wave quantum fluctuations during inflation, only the adiabatic fluctuation is important. We derive the relevant equations and compare with the conventional approach of the single-field approximation. As a result we show that this approximation is completely justified.

First, we consider inflation models in multifield configurations,  $\tilde{\varphi}$  and  $\tilde{\chi}_i$  ( $i = 1, \dots, N$ ). The Lagrangian density is given by

$$\mathcal{L} = -\frac{1}{2}(\partial_\mu \tilde{\varphi})^2 - \frac{1}{2}(\partial_\mu \tilde{\chi}_i)^2 - V(\tilde{\varphi}, \tilde{\chi}_i). \quad (4)$$

Here we assume that the field  $\tilde{\varphi}$  is light and the other fields  $\tilde{\chi}_i$  are heavy during inflation, that is,

$$|V_{,\tilde{\varphi}\tilde{\varphi}}| \ll H^2 \quad \text{and} \quad V_{,\tilde{\chi}_i\tilde{\chi}_i} \gg H^2, \quad (5)$$

where  $H \equiv \dot{a}/a$  is the Hubble parameter during inflation, a dot represents derivative with respect to the cosmic time  $t$ , and a comma represents partial derivative with respect to the corresponding field variables. Hereafter we set  $N = 1$  and denote  $\tilde{\chi}_1$  by  $\tilde{\chi}$  for simplicity. The extension to the case with  $N \geq 2$  is trivial.

We decompose the two fields to homogeneous modes and perturbations,

$$\begin{aligned} \tilde{\varphi}(t, \mathbf{x}) &= \varphi(t) + \delta\varphi(t, \mathbf{x}), \\ \tilde{\chi}(t, \mathbf{x}) &= \chi(t) + \delta\chi(t, \mathbf{x}). \end{aligned} \quad (6)$$

The perturbed metric around the flat Friedmann universe in the longitudinal gauge is taken as

$$ds^2 = -[1 + 2\Phi(t, \mathbf{x})] dt^2 + a(t)^2 [1 - 2\Phi(t, \mathbf{x})] \delta_{ij} dx^i dx^j \quad (7)$$

with  $\Phi(t, \mathbf{x})$  being the gravitational potential.

First of all, we consider the dynamics of the homogeneous modes  $\varphi(t)$  and  $\chi(t)$ . The equations of motion for the homogeneous modes are given by

$$\begin{aligned} \ddot{\varphi} + 3H\dot{\varphi} + V_{,\varphi} &= 0, \\ \ddot{\chi} + 3H\dot{\chi} + V_{,\chi} &= 0, \\ H^2 &= \frac{1}{3M_G^2} \left( \frac{1}{2}\dot{\varphi}^2 + \frac{1}{2}\dot{\chi}^2 + V \right). \end{aligned} \quad (8)$$

Since the mass of  $\chi$  is much larger than the Hubble parameter, the field  $\chi$  quickly relaxes to the minimum  $\chi^m = \chi^m(\varphi)$ , characterized by

$$V_{,\chi}(\varphi, \chi^m(\varphi)) = 0. \quad (9)$$

Since this relation holds for arbitrary  $\varphi$ , we find

$$\frac{dV_{,\chi}}{d\varphi}(\varphi, \chi^m(\varphi)) = V_{,\chi\varphi} + V_{,\chi\chi} \frac{d\chi^m}{d\varphi} = 0. \quad (10)$$

Defining the effective single-field potential  $V_s(\varphi)$  as  $V_s(\varphi) = V(\varphi, \chi = \chi^m(\varphi))$ , the equation of motion for  $\varphi$  reads

$$\ddot{\varphi} + 3H\dot{\varphi} + \frac{dV_s}{d\varphi} = 0, \quad (11)$$

where we have used

$$\frac{dV_s}{d\varphi}(\varphi, \chi^m(\varphi)) = \left. \frac{\partial V}{\partial \varphi} \right|_{\chi=\chi^m(\varphi)} + \frac{d\chi^m(\varphi)}{d\varphi} \left. \frac{\partial V}{\partial \chi} \right|_{\chi=\chi^m(\varphi)} = V_{,\varphi}(\varphi, \chi^m(\varphi)). \quad (12)$$

Thus, the dynamics of the homogeneous mode  $\varphi$  is completely determined by the reduced potential  $V_s(\varphi)$ . That is, we can look on this type of inflation as an effectively single-field inflation with the potential  $V_s(\varphi)$ , as far as the dynamics of the homogeneous mode is concerned.

Note that for successful inflation, the absolute magnitude of the effective mass squared of  $\varphi$  must be much smaller than the Hubble parameter squared, that is,

$$\left| \frac{d^2 V_s}{d\varphi^2} \right| = \left| V_{,\varphi\varphi} + 2 \frac{d\chi^m}{d\varphi} V_{,\varphi\chi} + \left( \frac{d\chi^m}{d\varphi} \right)^2 V_{,\chi\chi} \right| = \left| \frac{V_{,\varphi\varphi} V_{,\chi\chi} - V_{,\varphi\chi}^2}{V_{,\chi\chi}} \right| \ll H^2, \quad (13)$$

where we have used Eq. (10). One sufficient condition for this inequality is that the non-diagonal components of Hesse matrix of the potential  $V$  are not so large, that is,

$$V_{,\varphi\chi}^2 \lesssim |V_{,\varphi\varphi} V_{,\chi\chi}|. \quad (14)$$

Next, we consider perturbations of the field variables,  $\delta\varphi$  and  $\delta\chi$ , whose equations of motion are given by [17]

$$\begin{aligned} \ddot{\delta\varphi} + 3H\dot{\delta\varphi} - \frac{\nabla^2}{a^2} \delta\varphi + V_{,\varphi\varphi} \delta\varphi &= -2V_{,\varphi} \Phi + 4\dot{\varphi} \dot{\Phi} - V_{,\varphi\chi} \delta\chi, \\ \ddot{\delta\chi} + 3H\dot{\delta\chi} - \frac{\nabla^2}{a^2} \delta\chi + V_{,\chi\chi} \delta\chi &= -2V_{,\chi} \Phi + 4\dot{\chi} \dot{\Phi} - V_{,\chi\varphi} \delta\varphi. \end{aligned} \quad (15)$$

On the other hand, the evolution of the gravitational potential is governed by

$$\begin{aligned} \dot{\Phi} + H\Phi &= \frac{1}{2M_G^2} (\dot{\varphi} \delta\varphi + \dot{\chi} \delta\chi), \\ \ddot{\Phi} + H\dot{\Phi} + \frac{\nabla^2}{a^2} \Phi &= \frac{1}{M_G^2} (\dot{\varphi} \dot{\delta\varphi} + \dot{\chi} \dot{\delta\chi}). \end{aligned} \quad (16)$$

We are interested only in the adiabatic density fluctuations, which is characterized by the condition [7],

$$\frac{\delta\varphi}{\dot{\varphi}} = \frac{\delta\chi}{\dot{\chi}}, \quad (17)$$

or equivalently

$$\delta\chi = \frac{d\chi^m(\varphi)}{d\varphi}\delta\varphi, \quad (18)$$

where we have used  $\dot{\chi}/\dot{\varphi} = d\chi^m(\varphi)/d\varphi$ . Inserting the relation (10), the adiabatic condition (18) can be rewritten as

$$\delta\chi = \frac{d\chi^m}{d\varphi}\delta\varphi = -\frac{V_{,\chi\varphi}}{V_{,\chi\chi}}\delta\varphi. \quad (19)$$

Under this condition, the equation of the perturbation  $\delta\varphi$  becomes

$$\delta\ddot{\varphi} + 3H\delta\dot{\varphi} - \frac{\nabla^2}{a^2}\delta\varphi + \left(\frac{V_{,\varphi\varphi}V_{,\chi\chi} - V_{,\varphi\chi}^2}{V_{,\chi\chi}}\right)\delta\varphi = -2V_{,\varphi}\Phi + 4\dot{\varphi}\dot{\Phi}. \quad (20)$$

Here note that  $\delta\varphi$  is still light, that is, the absolute magnitude of the effective mass squared of  $\delta\varphi$  is also much smaller than  $H^2$  due to Eq. (13).

The equations of the gravitational potential are then written as

$$\dot{\Phi} + H\Phi = \frac{1}{2M_G^2} \left[ 1 + \left( \frac{d\chi^m}{d\varphi} \right)^2 \right] \dot{\varphi}\delta\varphi, \quad (21)$$

$$\ddot{\Phi} + H\dot{\Phi} + \frac{\nabla^2}{a^2}\Phi = \frac{1}{M_G^2} \left\{ \left[ 1 + \left( \frac{d\chi^m}{d\varphi} \right)^2 \right] \dot{\varphi}\delta\dot{\varphi} + \frac{d^2\chi^m}{d\varphi^2} \frac{d\chi^m}{d\varphi} \dot{\varphi}^2 \delta\varphi \right\}. \quad (22)$$

From these equations, the gravitational potential can be described by  $\delta\varphi$ ,

$$\begin{aligned} \left( \dot{H} - \frac{\nabla^2}{a^2} \right) \Phi &= \frac{1}{2M_G^2} (\ddot{\varphi}\delta\varphi - \dot{\varphi}\delta\dot{\varphi} + \ddot{\chi}\delta\chi - \dot{\chi}\delta\dot{\chi}), \\ &= \frac{1}{2M_G^2} (\ddot{\varphi}\delta\varphi - \dot{\varphi}\delta\dot{\varphi}) \left[ 1 + \left( \frac{d\chi^m}{d\varphi} \right)^2 \right]. \end{aligned} \quad (23)$$

Here  $\dot{H}$  is given by

$$\dot{H} = -\frac{1}{2M_G^2}(\dot{\varphi}^2 + \dot{\chi}^2) = -\frac{1}{2M_G^2}\dot{\varphi}^2 \left[ 1 + \left( \frac{d\chi^m}{d\varphi} \right)^2 \right]. \quad (24)$$

Therefore, in the long wave limit, the gravitational potential is given by

$$\Phi = \frac{d}{dt} \left( \frac{\delta\varphi}{\dot{\varphi}} \right). \quad (25)$$

Thus, adiabatic fluctuations produced in this inflation model are completely characterized by equations (11), (20), and (25).

Next, we investigate whether the above two equations can be reproduced by considering a single-field inflation model with the effective potential  $V_s(\varphi) = V(\varphi, \chi^m(\varphi))$ . For this purpose, let us consider the following Lagrangian density,

$$\mathcal{L} = -\frac{1}{2}(\partial_\mu \tilde{\varphi})^2 - V_s(\tilde{\varphi}). \quad (26)$$

The equation of motion for the homogeneous mode  $\varphi$  is given by the same equation (11). The equation of motion of  $\delta\varphi$  for this system is given by

$$\delta\ddot{\varphi} + 3H\delta\dot{\varphi} - \frac{\nabla^2}{a^2}\delta\varphi + \frac{d^2V_s}{d\varphi^2}\delta\varphi = -2\frac{dV_s}{d\varphi}\Phi + 4\dot{\varphi}\dot{\Phi}. \quad (27)$$

The derivative of the effective potential  $V_s(\varphi) = V(\varphi, \chi^m(\varphi))$  can be described by the partial derivative of the original potential,

$$\begin{aligned} \frac{dV_s}{d\varphi} &= V_{,\varphi} + V_{,\chi} \frac{d\chi^m}{d\varphi} = V_{,\varphi}, \\ \frac{d^2V_s}{d\varphi^2} &= V_{,\varphi\varphi} + 2\frac{d\chi^m}{d\varphi}V_{,\varphi\chi} + \left(\frac{d\chi^m}{d\varphi}\right)^2 V_{,\chi\chi} = \frac{V_{,\varphi\varphi}V_{,\chi\chi} - V_{,\varphi\chi}^2}{V_{,\chi\chi}}. \end{aligned} \quad (28)$$

Inserting this relation to Eq. (27) yields

$$\delta\ddot{\varphi} + 3H\delta\dot{\varphi} - \frac{\nabla^2}{a^2}\delta\varphi + \left(\frac{V_{,\varphi\varphi}V_{,\chi\chi} - V_{,\varphi\chi}^2}{V_{,\chi\chi}}\right)\delta\varphi = -2V_{,\varphi}\Phi + 4\dot{\varphi}\dot{\Phi}, \quad (29)$$

which completely coincides with Eq. (20).

The equations of the gravitational potential are written as

$$\dot{\Phi} + H\Phi = \frac{1}{2M_G^2}\dot{\varphi}\delta\varphi, \quad (30)$$

$$\ddot{\Phi} + H\dot{\Phi} + \frac{\nabla^2}{a^2}\Phi = \frac{1}{M_G^2}t\varphi\delta\dot{\varphi}. \quad (31)$$

From these equations, the gravitational potential can be described by  $\delta\varphi$ ,

$$\left(\dot{H} - \frac{\nabla^2}{a^2}\right)\Phi = \frac{1}{2M_G^2}(\ddot{\varphi}\delta\varphi - \dot{\varphi}\delta\dot{\varphi}). \quad (32)$$

Note that this equation is different from the equation (23) by the factor  $[1 + (d\chi^m/d\varphi)^2]$ . However, here  $\dot{H}$  is given by

$$\dot{H} = -\frac{1}{2M_G^2}\dot{\varphi}^2. \quad (33)$$

Therefore, in the long wave limit, the gravitational potential is given by

$$\Phi = \frac{d}{dt} \left( \frac{\delta\varphi}{\dot{\varphi}} \right), \quad (34)$$

which completely coincides with Eq. (25).

Thus, this type of inflation can be regarded as an effectively single-field inflation with the reduced potential  $V_s(\varphi)$  not only for the dynamics of the homogeneous mode but also for generation of adiabatic density fluctuations.

Finally we comment on the physical meaning of the adiabatic condition (17) in our system. The homogeneous mode of the heavy field is assumed to stay at its minimum during inflation, that is,

$$V_{,\chi}(\varphi, \chi) = 0, \quad \text{namely, } \chi = \chi^m(\varphi). \quad (35)$$

Under this condition, we can show that the adiabatic condition (17) is equivalent to the following requirement,

$$V_{,\chi}(\tilde{\varphi}, \tilde{\chi}) = 0, \quad \text{namely,} \quad \tilde{\chi} = \chi^m(\tilde{\varphi}). \quad (36)$$

That is,  $\tilde{\chi}$  is related to  $\tilde{\varphi}$  through the same function  $\chi^m$  as the case of the homogeneous mode and the perturbed fields also stay at the minimum of  $\chi$  for adiabatic fluctuations. This can be easily shown. First, we expand  $V_{,\chi}(\tilde{\varphi}, \tilde{\chi})$  as

$$V_{,\chi}(\tilde{\varphi}, \tilde{\chi}) = V_{,\chi}(\varphi + \delta\varphi, \chi + \delta\chi) = V_{,\chi\varphi}(\varphi, \chi)\delta\varphi + V_{,\chi\chi}(\varphi, \chi)\delta\chi = V_{,\chi\chi}(\varphi, \chi) \left( -\frac{d\chi^m}{d\varphi}\delta\varphi + \delta\chi \right), \quad (37)$$

where we have used the relation (10) in the last equation. Then, it is obvious that  $V_{,\chi}(\tilde{\varphi}, \tilde{\chi}) = 0$  is equivalent to

$$\delta\chi = \frac{d\chi^m}{d\varphi}\delta\varphi = \frac{\dot{\chi}}{\dot{\varphi}}\delta\varphi, \quad (38)$$

which is just the adiabatic condition (17).

To conclude, we have discussed primordial density fluctuations produced by an inflation model with multiple scalar fields in which only one field is light and the others are heavy. We have shown that adiabatic density fluctuations of such an inflation can be completely reproduced by looking on it as an effectively single-field model with the reduced potential obtained by inserting the minima of the heavy fields to the original potential.

## Acknowledgments

We are grateful to Robert H. Brandenberger for his useful comments. M.Y. is supported in part by the project of the Research Institute of Aoyama Gakuin University. J.Y. was partially supported by the JSPS Grant-in-Aid for Scientific Research No. 16340076.

- 
- [1] A.H. Guth, Phys. Rev. **D23**, 347 (1981); K. Sato, Mon. Not. R. astr. Soc. **195**, 467 (1981). For a review, see, for example, A.D. Linde, Particle Physics and Inflationary Cosmology, (Harwood, Chur, Switzerland, 1990).
  - [2] S.W. Hawking, Phys. Lett. **115B**, 295 (1982); A.A. Starobinsky, Phys. Lett. **117B**, 175 (1982); A.H. Guth and S-Y. Pi, Phys. Rev. Lett. **49**, 1110 (1982).
  - [3] C. L. Bennett *et al.*, Astrophys. J., Suppl. Ser. **148**, 1 (2003); **148**, 175 (2003).
  - [4] H. V. Peiris *et al.*, Astrophys. J., Suppl. Ser. **148**, 213 (2003).
  - [5] W. C. Jones *et al.*, astro-ph/0507494; F. Piacentini *et al.*, astro-ph/0507507; astro-ph/0507514.
  - [6] R. Bousso and J. Polchinski, J. High Energy Phys. **0006**, 006 (2000) S. Kachru, R. Kallosh, A. Linde, and S. P. Trivedi, Phys. Rev. D **68**, 046005 (2003), L. Susskind, hep-th/0302219.
  - [7] A. A. Starobinsky, JETP Lett. **42**, 152 (1985); D. Polarski and A. A. Starobinsky, Nucl. Phys. **B385**, 623 (1992); Phys. Rev. D **50**, 6123 (1994); A. A. Starobinsky and J. Yokoyama, Proc. of The Fourth Workshop on General Relativity and Gravitation, ed. by K. Nakao *et al.* (Kyoto University, 1994), p. 381 (gr-qc/9502002); V. F. Mukhanov and P. J. Steinhardt, Phys. Lett. B **422**, 52 (1998); T. Chiba, N. Sugiyama, and J. Yokoyama, Nucl. Phys. B **530**,



- 304 (1998); A. A. Starobinsky, S. Tsujikawa and J. Yokoyama, Nucl. Phys. B **610**, 383 (2001); S. G. Nibbelink and B. J. W. van Tent, Class. Quantum Grav. **19**, 613 (2002).
- [8] R. Easther, hep-th/0407042.
  - [9] A. D. Linde, Phys. Lett. B **259**, 38 (1991); Phys. Rev. D **49**, 748 (1994).
  - [10] M. Kawasaki, M. Yamaguchi, and Jun'ichi Yokoyama, Phys. Rev. D **68**, 023508 (2003); M. Yamaguchi and J. Yokoyama, Phys. Rev. D **68**, 123520 (2003).
  - [11] A. D. Linde, Phys. Lett. **129B**, 177 (1983).
  - [12] A. D. Linde, Phys. Lett. **108B**, 389 (1982); A. Albrecht and P. J. Steinhardt, Phys. Rev. Lett. **48**, 1220 (1982).
  - [13] A. Linde, Phys. Lett. B **327**, 208 (1994); A. Linde and D. Linde, Phys. Rev. D **50**, 2456 (1994); A. Vilenkin, Phys. Rev. Lett. **72**, 3137 (1994).
  - [14] G. Lazarides and C. Panagiotakopoulos, Phys. Rev. D **52**, R559 (1995); R. Jeannerot, S. Khalil, and, G. Lazarides, Phys. Lett. B **506**, 344 (2001).
  - [15] M. Yamaguchi and J. Yokoyama, Phys. Rev. D **70**, 023513 (2004).
  - [16] M. Kawasaki, T. Takayama, M. Yamaguchi, and J. Yokoyama, hep-ph/0605271.
  - [17] J. M. Bardeen, Phys. Rev. D **22**, 1882 (1980); H. Kodama and M. Sasaki, Prog. Theor. Phys. Suppl. **78**, 1 (1984); V.F. Mukhanov, H.A. Feldman, and R. H. Brandenberger, Phys. Rep. **215**, 203 (1992).